Claims

- (1.) An isolated nucleic acid molecule comprising a sequence substantially identical to SEQ ID NO:252.
- 2. \ The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule comprises the sequence shown in SEQ ID NO:252 5
 - An isolated nucleic acid molecule comprising a sequence substantially identical to SEQ ID NO;105.
 - The isolated nucleic acid molecule of claim 3, wherein said nucleic acid molecule comprises the sequence shown in SEQ ID NO:105.
 - An isolated nucleic acid molecule comprising a sequence substantially 5. identical to SEQ ID NO:106.
 - The isolated nucleic acid-molecule of claim 5, wherein said nucleic acid 6. molecule comprises the sequence shown in SEQ ID NO:106.
 - 7. A substantially pure polypeptide comprising an amino acid sequence that is substantially identical to the amino acid/sequence of SEQ ID NO:253.
 - 8. The substantially pure polypeptide of claim 7, wherein said amino acid sequence comprises the sequence shown in SEQ ID NO:253.
 - 9. A substantially pure polypeptide comprising an amino acid sequence that is substantially identical to the amino acid sequence of SEQ ID NO:107.
- 10. The substantially pure polypeptide of clarm 9, wherein said amino acid 20 sequence comprises the sequence shown in SEQ ID NO:107.

10

U إواث F 24

40

į.

20

- 11. A substantially pure polypeptide comprising an amino acid sequence that is substantially identical to the amino acid sequence of SEQ ID NO:108.
- 12. The substantially pure polypeptide of claim 11, wherein said amino acid sequence comprises the sequence shown in SEQ ID NO:108.
- A method for identifying a compound which is capable of decreasing 13. the expression of a pathogenic virulence factor, said method comprising the steps of:
 - (a) providing a pathogenic cell expressing a nucleic acid molecule of claim 1; and
 - (b) contacting said pathogenic cell with a candidate compound, a decrease in expression of said nucleic acid molecule following contact with said candidate compound identifying a compound which decreases the expression of a pathogenic virulence factor.
 - The method of claim 13, wherein said pathogenic cell infects a 14. mammal.
 - The method of claim 13, wherein said pathogenic cell infects a plant. 15.
 - A method for identifying a compound which is capable of decreasing 16. the expression of a pathogenic virulence factor, said method comprising the steps of:
 - (a) providing a pathogenic cell expressing a nucleic acid molecule of claim 3; and
 - (b) contacting said pathogenic cell with a candidate compound, a decrease in expression of said nucleic acid molecule following contact with said candidate compound identifying a compound which decreases the expression of a pathogenic virulence factor.
 - The method of claim 16, wherein said pathogenic cell infects a 17. mammal.

20

25

5

and

- The method of claim 16, wherein said pathogenic cell infects a plant. 18.
- A method for identifying a compound which is capable of decreasing 19. the expression of a pathogenic virulence factor, said method comprising the steps of:
 - (a) providing a pathogenic cell expressing a nucleic acid molecule of claim 5;
- (b) contacting said pathogenic cell with a candidate compound, a decrease in expression of said nucleic acid molecule following contact with said candidate compound identifying a compound which decreases the expression of a pathogenic virulence factor.
 - The method of claim 19, wherein said pathogenic cell infects a mammal. 20.
 - The method of claim 19, wherein said pathogenic cell infects a plant. 21.
- A method for identifying a compound which binds a polypeptide, said 22. method comprising the steps of:
- (a) contacting a candidate compound with a substantially pure polypeptide comprising an amino acid sequence of claim 7 under conditions that allow binding; and
 - (b) detecting binding of the candidate compound to the polypeptide.
- A method for identifying a compound which binds a polypeptide, said method comprising the steps of:
- (a) contacting a candidate compound with a substantially pure polypeptide comprising an amino acid sequence of claim 9 under conditions that allow binding; and
 - (b) detecting binding of the candidate compound to the polypeptide.
- 24. A method for identifying a compound which binds a polypeptide, said method comprising the steps of:
- (a) contacting a candidate compound with a substantially pure polypeptide comprising an amino acid sequence of claim 11 under conditions that allow binding; and

- (b) detecting binding of the candidate compound to the polypeptide.
- 25. A method of treating a pathogenic infection in mammal, said method comprising the steps of:
 - (a) identifying a mammal having a pathogenic infection; and
- 5 (b) administering to said mammal a therapeutically effective amount of a composition which inhibits the expression or activity of a polypeptide encoded by a nucleic acid molecule of claim 1 in said pathogen.
 - 26. A method of treating a pathogenic infection in mammal, said method comprising the steps of:
 - (a) identifying a mammal having a pathogenic infection; and
 - (b) administering to said mammal a therapeutically effective amount of a composition which inhibits the expression or activity of a polypeptide encoded by a nucleic acid molecule of claim 3 in said pathogen.
 - 27. The method of claim 26, wherein said pathogen is *Pseudomonas* aeruginosa.
 - 28. A method of treating a pathogenic infection in mammal, said method comprising the steps of:
 - (a) identifying a mammal having a pathogenic infection; and
 - (b) administering to said mammal a therapeutically effective amount of a composition which inhibits the expression or activity of a polypeptide encoded by a nucleic acid molecule of claim 5 in said pathogen.
 - 29. The method of claim 28, wherein said pathogen is *Pseudomonas* aeruginosa.
 - 30. A method of treating a pathogenic infection in a mammal, said method

comprising the steps of:

5

10

47

4

₩ 15

alle test atte

-

- (a) identifying a mammal having a pathogenic infection; and
- (b) administering to said mammal a therapeutically effective amount of a composition which binds and inhibits a polypeptide encoded by an amino acid sequence of claim 5.
- 31. The method of claim 30, wherein said pathogen is *Pseudomonas* aeruginosa.
- 32. A method of treating a pathogenic infection in a mammal, said method comprising the steps of:
 - (a) identifying a mammal having a pathogenic infection; and
- (b) administering to said mammal a therapeutically effective amount of a composition which binds and inhibits a polypeptide encoded by an amino acid sequence of claim 7.
- 33. The method of claim 32, wherein said pathogen infection is caused by *Pseudomonas aeruginosa*.
- 34. A method of treating a pathogenic infection in a mammal, said method comprising the steps of:
 - (a) identifying a mammal having a pathogenic infection; and
- (b) administering to said mammal a ther apeutically effective amount of a composition which binds and inhibits a polypeptide encoded by an amino acid sequence of claim 9.
 - 35. The method of claim 34, wherein said pathogen infection is caused by *Pseudomonas aeruginosa*.
 - 36. A method of treating a pathogenic infection in mammal, said method

10

4.74 4.74 4.74 4.74 4.74 4.74

44

I

15

-

20

- (a) identifying a\mammal having a pathogenic infection; and
- (b) administering to said mammal a therapeutically effective amount of a composition which binds and inhibits a substantially pure polypeptide comprising an amino acid sequence of claim 11.
- 37. The method of claim 36, wherein said pathogen infection is caused by *Pseudomonas aeruginosa*.
- 38. A method of identifying a compound which inhibits the virulence of a *Pseudomonas* cell, said method comprising the steps of:
 - (a) providing a *Pseudomonas* cell;
 - (b) contacting said cell with a candidate compound; and
- (c) detecting the presence of a phenazine, wherein a decrease in said phenazine relative to an untreated control cell is an indication that the compound inhibits the virulence of said *Pseudomonas* cell.
 - 39. The method of claim 38, wherein said cell is Pseudomonas aeruginosa.
- 40. The method of claim 38, wherein said phenazine is detected by spectroscopy.
 - 41. The method of claim 38, wherein said phenazine is a pyocyanin.
- 42. The method of claim 41, wherein said pyocyanin is detected by measuring the absorbance at 520 nm
 - 43. The method of claim 38, wherein said cell is present in a cell culture.

add A'

- 81 -